

Criss-cross reduction of the Maslov index and a proof of the Yoshida-Nicolaescu theorem

Booss-Bavnbek, Bernhelm

Publication date:
1999

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Booss-Bavnbek, B. (1999). *Criss-cross reduction of the Maslov index and a proof of the Yoshida-Nicolaescu theorem*. Roskilde Universitet. Tekster fra IMFUFA No. 199

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 373

1999

**Criss-Cross Reduction
of the Maslov Index
and a Proof of
the Yoshida-Nicolaescu Theorem**

**Bernhelm Booss-Bavnbek, Kenro Furutani,
and Nobukazu Otsuki**

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

IMFUFA, Roskilde University, P.O.Box 260, 4000 Roskilde, Denmark

Criss-Cross Reduction of the Maslov Index and a Proof of the Yoshida-Nicolaescu Theorem

By: Bernhelm Booss-Bavnbek, Kenro Furutani, and Nobukazu Otsuki

IMFUFA text nr. 373/99

15 pages

ISSN 01066242

ABSTRACT. We consider direct sum decompositions $\beta = \beta_- \dot{+} \beta_+$ and $L = L_- \dot{+} L_+$ of two symplectic Hilbert spaces by Lagrangian subspaces with dense embeddings $\beta_- \hookrightarrow L_-$ and $L_+ \hookrightarrow \beta_+$. We show that such criss-cross embeddings induce a continuous mapping between the Fredholm Lagrangian Grassmannians $\mathcal{FL}_{\beta_-}(\beta)$ and $\mathcal{FL}_{L_-}(L)$ which preserves the Maslov index for curves. This gives a slight generalization and a new proof of the Yoshida-Nicolaescu Spectral Flow Formula for families of Dirac operators over partitioned manifolds.

Criss-Cross Reduction of the Maslov Index and a Proof of the Yoshida-Nicolaescu Theorem

Bernhelm Booss-Bavnbek, Kenro Furutani, and Nobukazu Otsuki

ABSTRACT. We consider direct sum decompositions $\beta = \beta_- \dot{+} \beta_+$ and $L = L_- \dot{+} L_+$ of two symplectic Hilbert spaces by Lagrangian subspaces with dense embeddings $\beta_- \hookrightarrow L_-$ and $L_+ \hookrightarrow \beta_+$. We show that such criss-cross embeddings induce a continuous mapping between the Fredholm Lagrangian Grassmannians $\mathcal{FL}_{\beta_-}(\beta)$ and $\mathcal{FL}_{L_-}(L)$ which preserves the Maslov index for curves. This gives a slight generalization and a new proof of the Yoshida-Nicolaescu Spectral Flow Formula for families of Dirac operators over partitioned manifolds.

Introduction

As conjectured in [1], some spectral invariants of a closed partitioned manifold $M = M_0 \cup_{\Sigma} M_1$ with $M_0 \cap M_1 = \partial M_0 = \partial M_1 = \Sigma$ can be coded by the intersection geometry of the Cauchy data spaces along the partitioning hypersurface Σ of associated differential operators. The Yoshida-Nicolaescu Formula belongs to this program. It was proved in [16] in dimension 3 and subsequently generalized in [12] and modified by several authors (see e.g. [2], [3], [7], [8]). It expresses the spectral flow of a family of Dirac operators with the same principal symbol but continuously varying connections by the Maslov intersection index of the Cauchy data spaces.

A closer look at the presently available spectral flow formulas shows that they differ quite a bit in the kind and strength of the underlying assumptions and the substance of the claims made. It seems to us that our framework (based on standard functional analysis involving only elementary distribution theory) permits a wider applicability than rather deep means (like the symbolic calculus and approximation theory).

First of all, the Cauchy data spaces are treated in slightly different ways. On one side, the Cauchy data spaces are established as L^2 -closures of smooth

1991 *Mathematics Subject Classification.* Primary 58G25, Secondary 58G20.

Key words and phrases. Cauchy data spaces, elliptic boundary problems, families of Dirac operators, Lagrangian subspaces, Maslov index, spectral flow, symplectic functional analysis.

sections over the partitioning hypersurface Σ , coming from the restriction to the boundary of all smooth solutions over one of the parts M_j of the partitioned manifold M . In the case of the Dirac operator, this Cauchy data space can be represented as the range of the L^2 -extension of the pseudo-differential Calderón projection and established as a Lagrangian subspace of the symplectic Hilbert space $L^2(-\Sigma) \dot{+} L^2(\Sigma)$ (for the somewhat delicate details see e.g. [4], pp. 75–104).

On the other side, the Cauchy data spaces can be established as subspaces of the symplectic Hilbert space $\beta := D_{\max}/D_{\min}$ of natural boundary values, i.e. the boundary values of sections belonging to the maximal domain D_{\max} of the operator (so in [2], [3], see also [11]). One can embed β as a non-closed subspace into the distribution space $H^{-1/2}(-\Sigma) \dot{+} H^{-1/2}(\Sigma)$. This treatment of the Cauchy data spaces is independent of pseudo-differential analysis and has several conceptual and technical advantages: the arguments work for any symmetric elliptic differential operator of first order which satisfies a weak unique continuation property and admits an extension defining a self-adjoint Fredholm operator. Moreover, no product structures are required near the boundary. Finally, the space β is basic and most suitable for a short and general proof of the closedness and the Lagrangian property of the Cauchy data spaces and the continuity of the corresponding transformation from symmetric operators to Lagrangian subspaces. As observed already in [14], establishing this continuity is the crucial step in proving spectral flow formulas.

In [2], this β -program was carried out by standard functional analysis arguments and without assuming regularity at the endpoints or differentiability of the curve of operators, or other supplementary technical assumptions. The result was a quite general spectral flow formula, and the present paper is a continuation of [2]. We show, also by standard arguments, that an additional continuous transformation can be obtained, at least in the case when all metric structures are product near the separating (or non-separating) hypersurface, namely from the Cauchy data spaces of the β -theory into the ‘conventional’ L^2 -theory. This gives a slight generalization and a new proof of the Yoshida–Nicolaescu Formula.

In Section 1 we address a fairly general situation in symplectic functional analysis and prove a ‘criss-cross’ reduction theorem for the Maslov index. In Section 2 we give a condensed version of the General Spectral Flow Formula of [2]. In Section 3 we connect the two preceding sections and give a new proof of the Yoshida–Nicolaescu Formula. In the Appendix we correct an erroneous description in [2], §§ 1.1–1.2, and add a lemma to be inserted at the end of § 1.2 in [2].

1. Criss-Cross Reduction of the Maslov Index

Let β and L be symplectic Hilbert spaces with symplectic forms ω_β and ω_L , respectively. Let

$$(1.1) \quad \beta = \beta_- \dot{+} \beta_+ \quad \text{and} \quad L = L_- \dot{+} L_+$$

be direct sum decompositions by transversal (not necessarily orthogonal) pairs of Lagrangian subspaces. We assume that there exist continuous, injective mappings

$$(1.2) \quad i_- : \beta_- \longrightarrow L_- \quad \text{and} \quad i_+ : L_+ \longrightarrow \beta_+$$

with dense images and which are compatible with the symplectic structures, i.e.

$$(1.3) \quad \omega_L(i_-(x), a) = \omega_\beta(x, i_+(a)) \quad \text{for all } a \in L_+ \text{ and } x \in \beta_-.$$

Let λ_0 be a fixed Lagrangian subspace of β . We denote the Fredholm Lagrangian Grassmannian of λ_0 by $\mathcal{FL}_{\lambda_0}(\beta)$. It is the space of all Lagrangian subspaces of β which make a Fredholm pair with λ_0 . Its topology is defined by the operator norm of the projections. Its fundamental group is \mathbf{Z} , and the mapping of the loops in $\mathcal{FL}_{\lambda_0}(\beta)$ onto \mathbf{Z} is given by the Maslov index. It is an intersection index of the loop with the Maslov cycle

$$\mathcal{M}_{\lambda_0}(\beta) := \{\lambda \in \mathcal{FL}_{\lambda_0}(\beta) \mid \lambda \cap \lambda_0 \neq \{0\}\}.$$

By functional analytical means, the definition of the Maslov index can be extended unambiguously to all continuous curves in $\mathcal{FL}_{\lambda_0}(\beta)$ (see [2] and [3], inspired by [15]; we refer to [6] for various aspects of the Maslov index and also to [9], [10] for a cohomological treatment).

We are going to prove the following

THEOREM 1.1. *Under the assumptions (1.1), (1.2), and (1.3),*

(a) *we have a continuous mapping*

$$\tau : \mathcal{FL}_{\beta_-}(\beta) \longrightarrow \mathcal{FL}_{L_-}(L)$$

(b) *which maps the Maslov cycle of β_- into the Maslov cycle of L_- and*

(c) *preserves the Maslov index*

$$\text{mas}(\{\lambda_t\}_{t \in [0,1]}, \beta_-) = \text{mas}(\{\tau(\lambda_t)\}_{t \in [0,1]}, L_-)$$

for any continuous curve $[0, 1] \ni t \mapsto \lambda_t \in \mathcal{FL}_{\beta_-}(\beta)$.

We prove Theorem 1.1 in a series of small lemmata.

1.1. Definition of the mapping τ . We consider the direct sum

$$\mathcal{D} := \beta_+ \oplus L_- ,$$

where β and L are identified with subspaces of \mathcal{D} . Then we define the mapping τ simply by

$$(1.4) \quad \tau(\lambda) := \lambda \cap L \quad \text{for } \lambda \in \mathcal{FL}_{\beta_-}(\beta).$$

Clearly

$$(1.5) \quad \tau(\lambda) = \{i_-(a) + x \mid a \in \beta_-, x \in L_+ \text{ such that } i_+(x) + a \in \lambda\}.$$

To prove that $\text{range}(\tau) \subset \mathcal{FL}_{L_-}(L)$ and that τ is continuous, we introduce an alternative description of $\tau(\lambda)$ in terms of bounded operators associated to λ . For a given $\lambda \in \mathcal{FL}_{\beta_-}(\beta)$, we fix a direct sum decomposition

$$\lambda = (\lambda \cap \beta_-) \dot{+} \mu$$

with a suitable closed μ . Let $\pi_+ : \beta \rightarrow \beta_+$ denote the projection along β_- . Then to claim that (λ, β_-) is a Fredholm pair is equivalent to claiming that the projection $\pi_\lambda := \pi_+|_\lambda : \lambda \rightarrow \beta_+$ is a Fredholm operator. So we deduce that $F_\lambda := \pi_+(\lambda) = \pi_\lambda(\mu)$ is closed. For later use we notice that

$$(1.6) \quad \dim \beta_+ / F_\lambda = \dim(\beta_+ \dot{+} \beta_-) / (\lambda \dot{+} \beta_-) = \dim \lambda \cap \beta_- < +\infty.$$

By the injectivity of $\pi_+|_\mu$ we can write μ as the graph of a uniquely determined bounded operator

$$f_\mu : F_\lambda \longrightarrow \beta_-.$$

Then we rewrite

$$(1.7) \quad \tau(\lambda) = i_-(\lambda \cap \beta_-) + \text{graph}(\varphi_\lambda),$$

where

$$\begin{array}{ccc} \varphi_\lambda : i_+^{-1}(F_\lambda) & \longrightarrow & L_- \\ x & \mapsto & i_- \circ f_\mu \circ i_+(x) \end{array}$$

Since φ_λ is bounded and its domain is closed in L , its graph is also closed in L , and so is $\tau(\lambda)$ which differs from $\text{graph}(\varphi_\lambda)$ only by a space of finite dimension. But moreover:

LEMMA 1.2. *For each $\lambda \in \mathcal{FL}_{\beta_-}(\beta)$, we have $\tau(\lambda) \in \mathcal{FL}_{L_-}(L)$.*

PROOF. In the exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_\lambda & \hookrightarrow & \beta_+ & \xrightarrow{p} & \beta_+ / F_\lambda \longrightarrow 0 \\ & & & & \uparrow i_+ & & \\ & & & & L_+ & & \end{array}$$

the range of i_+ is dense in β_+ and $\dim \beta_+ / F_\lambda < +\infty$. So, the map $p \circ i_+$ is surjective. We therefore have

$$(1.8) \quad \beta_+ = F_\lambda + i_+(L_+)$$

and $\ker(p \circ i_+) = i_+^{-1}(F_\lambda)$. It follows that

$$(1.9) \quad L_+ / i_+^{-1}(F_\lambda) \cong (F_\lambda + i_+(L_+)) / F_\lambda = \beta_+ / F_\lambda.$$

This implies that the closed subspace $i_+^{-1}(F_\lambda)$ of L_+ is of finite codimension.

Now let $x, y \in i_+^{-1}(F_\lambda)$ and set $a := i_+(x)$ and $b := i_+(y)$. Then

$$\begin{aligned}\omega_L(x + \varphi_\lambda(x), y + \varphi_\lambda(y)) &= \omega_L(x, \varphi_\lambda(y)) + \omega_L(\varphi_\lambda(x), y) \\ &= \omega_\beta(a, f_\mu(b)) + \omega_\beta(f_\mu(a), b) \\ &= \omega_\beta(a + f_\mu(a), b + f_\mu(b)) = 0\end{aligned}$$

by the compatibility condition (1.3) and the isotropy of μ . So $\text{graph } \varphi_\lambda$ is isotropic. Moreover, we have

$$\dim(\text{graph}(\varphi_\lambda)^0 / \text{graph}(\varphi_\lambda)) = 2 \dim(L_+ / i_+^{-1}(F_\lambda)) = 2 \dim(\beta_+ / F_\lambda).$$

Here $\text{graph}(\varphi_\lambda)^0$ denotes the annihilator of $\text{graph}(\varphi_\lambda)$ with respect to the symplectic form ω_L . Clearly,

$$i_-(\lambda \cap \beta_-) \subset \text{graph}(\varphi_\lambda)^0 \quad \text{and} \quad i_-(\lambda \cap \beta_-) \cap \text{graph}(\varphi_\lambda) = \{0\}.$$

Further, we have

$$\dim i_-(\lambda \cap \beta_-) = \dim(\lambda \cap \beta_-) = \dim(\beta_+ / F_\lambda),$$

since the Fredholm index of the Lagrangian pair (λ, β_-) vanishes. From this dimension examination we see that adding $i_-(\lambda \cap \beta_-)$ to $\text{graph}(\varphi_\lambda)$ makes $\tau(\lambda)$ a Lagrangian subspace of L . It also follows that $\tau(\lambda) \cap L_- = i_-(\lambda \cap \beta_-)$ is of finite dimension and that $L_- + \tau(\lambda) = L_- + \text{graph}(\varphi_\lambda)$ is of finite codimension, hence $(\tau(\lambda), L_-)$ is a Fredholm pair in L . \square

REMARK 1.3. From equation (1.7) it particularly follows that

$$\tau(\mathcal{M}_{\beta_-}(\beta)) \subset \mathcal{M}_{L_-}(L).$$

1.2. The continuity of τ . We fix a closed $W \subset L_-$ with $\dim L_- / W < +\infty$ and choose a Lagrangian subspace θ of β with $\theta \pitchfork \beta_+$ and $W_\beta \subset \theta \subset W_\beta^0$, where $W_\beta := i_-^{-1}(W)$. Here “ \pitchfork ” means that the two subspaces intersect transversally. We notice that

$$W^0 = L_- \dot{+} (L_+ \cap W^0)$$

and, correspondingly,

$$W_\beta^0 = \beta_- \dot{+} (\beta_+ \cap W_\beta^0).$$

Next we exploit that the injection $i_- : \beta_- \rightarrow L_-$ has a dense range. Similar to the short exact sequence at the beginning of the proof of Lemma 1.2, we have

$$\begin{array}{ccccccc} 0 & \longrightarrow & W & \hookrightarrow & L_- & \xrightarrow{q} & L_- / W \longrightarrow 0 \\ & & & & \uparrow i_- & & \\ & & & & \beta_- & & \end{array}$$

We deduce, as above,

$$L_- = W + i_-(\beta_-) \quad \text{and} \quad \dim(\beta_- / W_\beta) = \dim(L_- / W).$$

Moreover, we have

$$(1.10) \quad i_+(W^0 \cap L_+) = W_\beta^0 \cap \beta_+.$$

To show the inclusion " \subset ", we consider any $x \in W^0 \cap L_+$. Then $\omega_L(x, y) = 0$ for all $y \in W$, and hence also for $y = i_-(z)$ for any $z \in W_\beta = i_-^{-1}(W)$. Now, by (1.3),

$$0 = \omega_L(x, i_-(z)) = \omega_\beta(i_+(x), z).$$

So,

$$i_+(x) \in W_\beta^0 \cap \beta_+.$$

We deduce the identity " $=$ " by the following dimension examination:

$$2 \dim(W^0 \cap L_+) = \dim(W^0/W) = \dim(W_\beta^0/W_\beta) = 2 \dim(W_\beta^0 \cap \beta_+).$$

Equation (1.10) permits us to rewrite

$$W_\beta^0 = i_+(L_+ \cap W^0) \dot{+} \beta_-$$

and to define

$$\begin{aligned} i_W : \quad W_\beta^0 &\longrightarrow W^0 \\ i_+(x) + z &\longmapsto x + i_-(z) \end{aligned}$$

for $x \in L_+ \cap W^0$ and $z \in \beta_-$. We obtain a new splitting of the symplectic Hilbert spaces:

PROPOSITION 1.4. *The space $\eta := i_W(\theta) + W$ is a Lagrangian subspace of L and the mapping $i_W|_\theta : \theta \rightarrow \eta$ has a dense image. Further, we have new direct sum decompositions*

$$\beta = \beta_+ \dot{+} \theta \quad \text{and} \quad L = L_+ \dot{+} \eta$$

which are compatible with regard to the symplectic forms ω_L and ω_β (similar to (1.3)).

PROOF. Clearly

$$W \subset \eta = i_W(\theta) + W \subset W^0 \quad \text{and} \quad \dim W^0/W < +\infty,$$

hence η is also closed and isotropic. We notice that the mapping $q \circ i_W|_\theta$ is surjective with $\ker q \circ i_W|_\theta = W_\beta$, where $q : \eta \rightarrow \eta/W$ denotes the projection. So,

$$W_\beta^0/W_\beta \supset \theta/W_\beta \cong \eta/W \subset W^0/W.$$

Then, from the dimension examination

$$\dim(W_\beta^0/W_\beta) = \dim(W^0/W),$$

it follows that η is a Lagrangian subspace of L .

To see that the range of $i_W|_\theta$ is dense in η we recall that the mapping $i_W : W_\beta^0 \rightarrow W^0$ is an isomorphism $W_\beta^0 \cap \beta_+ \cong W^0 \cap L_+$ on the first finite-dimensional component, and it is equal to the dense embedding $i_- : \beta_- \rightarrow L_-$ on the second component, hence its restriction to θ has a dense range in η .

The new direct sum decompositions and the compatibility of the symplectic forms follow at once. \square

The preceding proposition permits a further simplification of the graph representation of $\tau(\lambda)$, obtained in (1.7):

COROLLARY 1.5. *For any $\lambda \in \mathcal{FL}_{\beta_-}(\beta)$ with $\lambda \pitchfork \theta$, we have*

$$\tau(\lambda) = \text{graph}(i_W \circ f_\lambda \circ i_+),$$

where $f_\lambda : \beta_+ \rightarrow \theta$ such that $\text{graph } f_\lambda = \lambda$.

Now we can prove the continuity of τ and the invariance of the Maslov index under τ . We set

$$\mathcal{FL}_{W_\beta}(\beta)^{(0)} := \{\mu \in \mathcal{FL}_{W_\beta}(\beta) \mid \mu \cap W_\beta = \{0\}\}.$$

Then we have an open covering

$$\bigcup_W \mathcal{FL}_{W_\beta}(\beta)^{(0)} = \mathcal{FL}_{\beta_-}(\beta),$$

where the union is taken over all closed subspaces $W \subset L_-$ of finite codimension. It follows that $\tau(\mathcal{FL}_{W_\beta}(\beta)^{(0)}) \subset \mathcal{FL}_W(L)^{(0)}$. Thus, we have established restrictions

$$\tau_W : \mathcal{FL}_{W_\beta}(\beta)^{(0)} \longrightarrow \mathcal{FL}_W(L)^{(0)}$$

of τ for each W .

We fix a W . We denote the space of Lagrangian subspaces in a finite-dimensional symplectic space by $\text{Lag}(\cdot)$ and define the reduction map

$$\begin{aligned} \rho_{W_\beta} : \mathcal{FL}_{W_\beta}(\beta)^{(0)} &\longrightarrow \text{Lag}(W_\beta^0/W_\beta) \\ \lambda &\longmapsto (\lambda \cap W_\beta^0 + W_\beta) / W_\beta \end{aligned}$$

Then, for each Lagrangian subspace θ in β with $\theta \supset W_\beta$ and $\theta \pitchfork \beta_+$ (as before), the set

$$U(W_\beta, \theta) := \{\Lambda \in \text{Lag}(W_\beta^0/W_\beta) \mid \Lambda \pitchfork \theta/W_\beta\}$$

is open in $\text{Lag}(W_\beta^0/W_\beta)$ and we have an open covering

$$\bigcup_{\theta \pitchfork \beta_+} \rho_{W_\beta}^{-1}(U(W_\beta, \theta)) = \mathcal{FL}_{W_\beta}(\beta)^{(0)}.$$

By Corollary 1.5, the mapping τ is continuous on each $\rho_{W_\beta}^{-1}(U(W_\beta, \theta))$, and so it is continuous on the whole $\mathcal{FL}_{\beta_-}(\beta)$. This concludes the proof of (a) of Theorem 1.1.

Moreover, we have

$$\tau(\rho_{W_\beta}^{-1}(U(W_\beta, \theta))) \subset \rho_W^{-1}(U(W, \tau_W(\theta))),$$

where the reduction map $\rho_W : \mathcal{FL}_W(L)^{(0)} \rightarrow \text{Lag}(W^0/W)$ is defined correspondingly to ρ_{W_β} . We denote by

$$\overline{\tau}_W : \text{Lag}(W_\beta^0/W_\beta) \longrightarrow \text{Lag}(W^0/W)$$

the mapping naturally induced by τ_W . Then we have the following finite-dimensional symplectic reduction:

PROPOSITION 1.6. *The diagram*

$$(1.11) \quad \begin{array}{ccccc} \mathcal{FL}_{\beta_-}(\beta) & \xleftarrow{\quad} & \mathcal{FL}_{W_\beta}(\beta)^{(0)} & \xrightarrow{\rho_{W_\beta}} & \text{Lag}(W_\beta^0/W_\beta) \\ \tau \downarrow & & \tau_W \downarrow & & \downarrow \cong \overline{\tau_W} \\ \mathcal{FL}_{L_-}(L) & \xleftarrow{\quad} & \mathcal{FL}_W(L)^{(0)} & \xrightarrow{\rho_W} & \text{Lag}(W^0/W) \end{array}$$

is commutative.

For any continuous curve $\{\lambda_t\}_{t \in [0,1]} \in \mathcal{FL}_{\beta_-}(\beta)$, we can find a closed subspace $W \subset L_-$ of finite codimension such that the whole curve is contained in $\mathcal{FL}_{W_\beta}(\beta)^{(0)}$, respectively $\{\tau(\lambda_t)\}_{t \in [0,1]}$ is contained in $\mathcal{FL}_W(L)^{(0)}$. By finite symplectic reduction we obtain at once:

COROLLARY 1.7. *The Maslov index coincides under the transformations τ , τ_W , and $\overline{\tau_W}$ for loops and also for paths.*

NOTE. For paths, the Maslov index depends on the choice of the Maslov cycle. In $\text{Lag}(W^0/W)$, the Maslov index is taken with respect to the Maslov cycle $\mathcal{M}_{L_-/W}(W^0/W)$. Correspondingly, the Maslov index in $\text{Lag}(W_\beta^0/W_\beta)$ is taken with respect to the Maslov cycle $\mathcal{M}_{\beta_-/W_\beta}(W_\beta^0/W_\beta)$. Note that the specified Maslov cycle in $\text{Lag}(W_\beta^0/W_\beta)$ is mapped onto the specified Maslov cycle in $\text{Lag}(W^0/W)$ by $\overline{\tau_W}$.

Proposition 1.6 and Corollary 1.7 together with Remark 1.3 give the proof of (b) and (c) of Theorem 1.1.

2. The General Spectral Flow Formula

Let \mathcal{H} be a real separable Hilbert space and A an (unbounded) closed symmetric operator defined on the domain D_{\min} which is supposed to be dense in \mathcal{H} . Let A^* denote its adjoint operator with domain D_{\max} . We have that $A^*|_{D_{\min}} = A$ and that A^* is the maximal closed extension of A in \mathcal{H} .

We form the space β of *natural boundary values* with the *natural trace map* γ in the following way:

$$\begin{array}{ccc} D_{\max} & \xrightarrow{\gamma} & D_{\max}/D_{\min} =: \beta \\ x & \mapsto & \gamma(x) = [x] := x + D_{\min} \end{array}$$

The space β becomes a symplectic Hilbert space with the scalar product induced by the graph norm

$$(2.1) \quad (x, y)_\beta := (x, y) + (Ax, Ay)$$

and the symplectic form given by Green's form

$$(2.2) \quad \omega([x], [y]) := (Ax, y) - (x, Ay) \quad \text{for } [x], [y] \in \beta.$$

We define the *natural Cauchy data space* $\Lambda := \gamma(\ker A^*)$. It is a Lagrangian subspace of β under the assumption that A admits at least one self-adjoint Fredholm extension A_D . Actually, we shall assume a little more, namely that

A has a self-adjoint extension A_D with compact resolvent. Then $(\Lambda, \gamma(D))$ is a Fredholm pair of subspaces of β , i.e. $\Lambda \in \mathcal{FL}_{\gamma(D)}(\beta)$.

We consider a continuous curve $\{C_t\}_{t \in [0,1]}$ in the space of bounded self-adjoint operators on \mathcal{H} and assume that the operators $A^* + C_t - r$ have no 'inner solutions', i.e. satisfy the *weak unique continuation property* (UCP)

$$\ker(A^* + C_t - r) \cap D_{\min} = \{0\}$$

for $t \in [0, 1]$ and $|r| < \varepsilon_0$ with $\varepsilon_0 > 0$.

Clearly, the domains D_{\max} and D_{\min} are unchanged by the perturbation C_t for any t . So, β does not depend on the parameter t . Moreover, the symplectic form ω is invariantly defined on β and so also independent of t . It follows (see [2], Theorem 3.9) that the curve $\{\Lambda_t := \gamma(\ker(A^* + C_t))\}$ is continuous in $\mathcal{FL}_{\gamma(D)}(\beta)$.

Given this, the family $\{A_D + C_t\}$ can be considered at the same time in the spectral theory of self-adjoint operators, defining a spectral flow, and in the symplectic category, defining a Maslov index. Under the preceding assumptions, the main result obtainable at that level is the following general spectral flow formula (proved in [2], Theorem 5.1):

THEOREM 2.1. *Let A_D be a self-adjoint extension of A with compact resolvent and let $\{A_D + C_t\}$ be a family satisfying the UCP assumption. Then*

$$\text{sf}\{A_D + C_t\} = \text{mas}(\{\Lambda_t\}, \gamma(D)).$$

3. A Proof of the Yoshida–Nicolaescu Formula

Let M be a closed connected smooth Riemannian manifold and let $\Sigma \subset M$ be a hypersurface. Let

$$\{A_t := A_0 + C_t : C^\infty(M; S) \longrightarrow C^\infty(M; S)\}_{0 \leq t \leq 1}$$

be a continuous family of symmetric elliptic differential operators of first order with the same principal symbol, acting on sections of a real bundle S over M . The variation of the fixed operator A_0 is given by a continuous family $\{C_t\}_{t \in [0,1]}$ of smooth symmetric bundle homomorphisms. We assume that the normal bundle of the hypersurface Σ is orientable and that all metric structures of M and S are *product* in a collar neighbourhood $\mathcal{N} = (-1, 1) \times \Sigma$ of Σ . We have

$$(3.1) \quad A_t = \sigma \left(\frac{\partial}{\partial \tau} + B_t \right) \text{ on } \mathcal{N},$$

where τ denotes the normal coordinate, σ is orthogonal (assuming $\sigma^2 = -\text{Id}$), $\sigma B_t = -B_t \sigma$, and B_t is a self-adjoint elliptic differential operator over Σ , called the *tangential operator*. Here the point of the product structure is that then σ and B_t do not depend on the normal variable.

We cut the manifold at Σ and attach a copy of Σ to each side. So, we obtain a new manifold $M_\#$ with boundary $\Sigma_0 \sqcup \Sigma_1 = (-\Sigma) \sqcup \Sigma$. Then the β -space of

M_{\sharp} , being a $C^{\infty}(\partial M_{\sharp})$ -module, splits according to the connected components of ∂M_{\sharp} and we obtain two symplectic Hilbert spaces

$$\beta := \beta_0 \dot{+} \beta_1 \quad \text{and} \quad L := L_0 \dot{+} L_1$$

with

$$\gamma = \gamma_0 \oplus \gamma_1 : D_{\max}(M_{\sharp}) \longrightarrow \beta_0 \dot{+} \beta_1$$

and $L_j = L^2(\pm\Sigma; S|_{\pm\Sigma})$ for $j = 0, 1$. We note that L_0 and L_1 are naturally identified as Hilbert spaces, but their symplectic forms have opposite signs. In β we have a Lagrangian subspace

$$(3.2) \quad \Delta_{\beta} := \{(x, x) \in \beta_0 \dot{+} \beta_1 \mid \gamma_0(s) = x = \gamma_1(s) \text{ for } s \in H^1(M)\},$$

where $H^1(M)$ denotes the first order Sobolev space. Since M is closed, the operator A_t on $H^1(M)$ is a self-adjoint Fredholm operator. It can be identified with the operator $A_{t,D}^{\sharp} = A_{0,D}^{\sharp} + C_t$ over the new manifold M_{\sharp} with domain

$$D := \{s \in D_{\max}(M_{\sharp}) \mid \gamma_0(s) = \gamma_1(s)\},$$

i.e. it can be considered as a global self-adjoint boundary problem with $\gamma(D) = \Delta_{\beta}$. (Actually, D coincides with $H^1(M)$).

We set $\beta_- := \Delta_{\beta}$. In L , we have a Lagrangian subspace

$$(3.3) \quad L_- := \text{the diagonal of } L^2(-\Sigma; S|_{-\Sigma}) \dot{+} L^2(\Sigma; S|_{\Sigma}).$$

Clearly, the embedding $\beta_- \hookrightarrow L_-$ is bounded and dense.

To define suitable complementary Lagrangian subspaces β_+ and L_+ , we must fall back on a spectral resolution $\{\varphi_k, \lambda_k\}$ of $L^2(\Sigma)$ by eigensections of B_0 . (Here and in the following we do not mention the bundle S). For simplicity, we assume $\ker B_0 = \{0\}$. Otherwise we must decompose the finite-dimensional symplectic vector space $\ker B_0$ into two Lagrangian subspaces and add these spaces to the half parts defined by the spectral cut at 0.

Following [3], Proposition 7.15 (for related results see also [5] and [11]), we decompose

$$\beta_0 = \beta_-^0 \dot{+} \beta_+^0 \quad \text{and} \quad \beta_1 = \beta_-^1 \dot{+} \beta_+^1,$$

where

$$\beta_-^1 := \overline{[\{\varphi_k\}_{k < 0}]}^{H^{\frac{1}{2}}(\Sigma)} \quad \text{and} \quad \beta_+^1 := \overline{[\{\varphi_k\}_{k > 0}]}^{H^{-\frac{1}{2}}(\Sigma)}$$

and

$$\beta_-^0 := \overline{[\{\varphi_k\}_{k < 0}]}^{H^{-\frac{1}{2}}(\Sigma)} \quad \text{and} \quad \beta_+^0 := \overline{[\{\varphi_k\}_{k > 0}]}^{H^{\frac{1}{2}}(\Sigma)}.$$

In a similar way, we decompose

$$L_0 = L_-^0 \dot{+} L_+^0 \quad \text{and} \quad L_1 = L_-^1 \dot{+} L_+^1$$

with

$$L_-^1 := \overline{[\{\varphi_k\}_{k < 0}]}^{L^2(\Sigma)} \quad \text{and} \quad L_+^1 := \overline{[\{\varphi_k\}_{k > 0}]}^{L^2(\Sigma)}$$

and

$$L_-^0 := \overline{[\{\varphi_k\}_{k<0}]^{L^2(\Sigma)}} \quad \text{and} \quad L_+^0 := \overline{[\{\varphi_k\}_{k>0}]^{L^2(\Sigma)}}.$$

Rewriting

$$(3.4) \quad \beta = \beta_0 \dot{+} \beta_1 = \beta_-^0 \dot{+} \beta_+^0 \dot{+} \beta_-^1 \dot{+} \beta_+^1,$$

we obtain

$$\beta_- = \Delta_\beta = \{(a, b, a, b) \mid (a, b) = \gamma_0(s) = \gamma_1(s) \text{ with } s \in H^1(M)\}.$$

Correspondingly, we define

$$\beta_+ := \{(x, 0, 0, y) \mid x \in \beta_-^0, y \in \beta_+^1\}$$

and

$$L_+ := \{(u, 0, 0, v) \mid u \in L_-^0, v \in L_+^1\}.$$

Clearly, we have $L = L_- \dot{+} L_+$, a dense bounded embedding $L_+ \hookrightarrow \beta_+$, and $\Delta_\beta \cap \beta_+ = \{0\}$. From the decomposition (3.4) we have also that $\Delta_\beta + \beta_+ = \beta$. Then, as explained above in Section 2, the β -theory gives the continuity of the family of Cauchy data spaces $\{\Lambda_t\}$ (of the continuous operator family $\{A_t\}$, considered over $M_\#$). They are all Lagrangian subspaces of β and make Fredholm pairs with $\beta_- = \Delta_\beta = \gamma(H^1(M))$. From the General Spectral Flow Formula (here Theorem 2.1) and by the criss-cross reduction of the Maslov index (Theorem 1.1), we obtain

$$(3.5) \quad \text{sf}\{A_0 + C_t\} = \text{sf}\{A_{0,D}^\# + C_t\}$$

$$(3.6) \quad \stackrel{\text{Th.2.1}}{=} \text{mas}(\{\Lambda_t\}, \Delta_\beta)$$

$$(3.7) \quad \stackrel{\text{Th.1.1}}{=} \text{mas}(\{\Lambda_t \cap L\}, L_-).$$

Note that here

$$\Lambda_t \cap L = \{s|_{(-\Sigma) \cup \Sigma} \mid s \in H^{\frac{1}{2}}(M_\#) \text{ and } A_t(s) = 0 \text{ in } M_\# \setminus \partial M_\#\},$$

i.e. it coincides with the 'conventional' L^2 -definition of the Cauchy data spaces.

In the particular case of a *partitioned* manifold

$$M = M_0 \cup_\Sigma M_1 \quad \text{with } \Sigma = \partial M_0 = \partial M_1 = M_0 \cap M_1,$$

not only β and L split but also the Cauchy data spaces split

$$\Lambda_t = \Lambda_t^0 \dot{+} \Lambda_t^1,$$

according to the splitting of M into two parts. So, we obtain from (3.7) the true Yoshida–Nicolaescu Formula, though without any assumptions about the regularity at the endpoints or the differentiability of the curve:

THEOREM 3.1.

$$\begin{aligned} \text{sf}\{A_0 + C_t\} &= \text{mas}(\{\Lambda_t^0 \cap L^2(-\Sigma) \dot{+} \Lambda_t^1 \cap L^2(\Sigma)\}, L_-) \\ &=: \text{mas}(\{\Lambda_t^0 \cap L^2(-\Sigma)\}, \{\Lambda_t^1 \cap L^2(\Sigma)\}), \end{aligned}$$

where the last expression is just a rewriting of the Maslov index of Fredholm pairs of two curves.

REMARK 3.2. (a) We notice that the subspaces β_- and L_- are defined independently of a reference tangential operator, here B_0 , but the choice of any other reference operator B_t would have given a different decomposition, though not a different result.

(b) We would like to emphasize that the Yoshida–Nicolaescu formulas in the literature always assume product structures near Σ , whereas the general spectral flow formulas, as proved in [2] and expressed in [3] in $\beta \subset H^{-1/2}(\Sigma)$, do not require product structures near Σ . However, we also need product structures near Σ , to apply our criss-cross reduction theorem and to transform the general spectral flow formulas, expressed in ‘natural’ distribution spaces, into ‘conventional’ L^2 -formulas. Possibly, the criss-cross reduction theorem may provide one explanation for the need of product structures for L^2 -formulas.

(c) We also want to point to a misprint in [3], p. 74 (after Equation (7.13)), where it must read that ‘ $\gamma(S) \cap L^2(\widehat{\Sigma})$ is not closed in $L^2(\widehat{\Sigma})$ ’ instead of ‘ $\gamma(S) \cap L^2(\Sigma)$ is not closed in $L^2(\Sigma)$ ’.

Appendix A. Addendum and Corrections to [2]

Let \mathcal{L} denote the space of all Lagrangian subspaces of a real separable symplectic Hilbert space \mathcal{H} , and let $\mathcal{L}^{\mathbb{C}}$ denote the space of all complex Lagrangian subspaces of $\mathcal{H} \otimes \mathbb{C}$. As correctly observed in [2], the full group $\mathcal{U}(\mathcal{H})$ of unitary operators of \mathcal{H} does *not* act on $\mathcal{FL}_{\lambda_0}(\mathcal{H})$, but the reduced group $\mathcal{U}_c(\mathcal{H})$ does. It consists of unitary operators of the form $\text{Id} + K$, where K is a compact operator. In [2], below on p. 5, we claimed that the reduced group acts transitively on $\mathcal{FL}_{\lambda_0}(\mathcal{H})$, i.e. the mapping

$$\begin{array}{ccc} \rho: & \mathcal{U}_c(\mathcal{H}) & \longrightarrow \mathcal{FL}_{\lambda_0}(\mathcal{H}) \\ & U & \mapsto U(\lambda_0^\perp) \end{array}$$

is surjective. This claim is erroneous, since the difference of the orthogonal projections onto a Fredholm pair of closed subspaces of \mathcal{H} need not be of the form $\text{Id} + K$, but can become an arbitrary Fredholm operator as proved in [3], Appendix.

However, if we replace $\mathcal{U}_c(\mathcal{H})$ by $\mathcal{U}(\mathcal{H})$, $\mathcal{FL}_{\lambda_0}(\mathcal{H})$ by \mathcal{L} , $\mathcal{U}_c(\mathcal{H} \otimes \mathbb{C})$ by $\mathcal{U}(\mathcal{H} \otimes \mathbb{C})$, and $\mathcal{FL}_{\lambda_0 \otimes \mathbb{C}}^{\mathbb{C}}(\mathcal{H} \otimes \mathbb{C})$ by $\mathcal{L}^{\mathbb{C}}$, we have the same results as in [2], §§ 1.1–1.2:

The group $\mathcal{U}(\mathcal{H})$ (resp. $\mathcal{U}(\mathcal{H} \otimes \mathbb{C})$) acts transitively on \mathcal{L} (resp. on $\mathcal{L}^{\mathbb{C}}$). So let

$$(A.1) \quad \begin{array}{ccc} \rho: & \mathcal{U}(\mathcal{H}) & \longrightarrow \mathcal{L} \\ & U & \mapsto U(\lambda_0^\perp) \end{array}$$

and

$$(A.2) \quad \begin{array}{ccc} \rho^{\mathbb{C}}: & \mathcal{U}(\mathcal{H} \otimes \mathbb{C}) & \longrightarrow \mathcal{L}^{\mathbb{C}} \\ & g & \mapsto g(\lambda_0^\perp \otimes \mathbb{C}) \end{array}$$

denote the mappings defined by these actions. We obtain a commutative diagram

$$(A.3) \quad \begin{array}{ccc} \mathcal{U}(\mathcal{H}) & \xrightarrow{\tilde{\tau}} & \mathcal{U}(\mathcal{H} \otimes \mathbb{C}) \\ \downarrow \rho & & \downarrow \rho^{\mathbb{C}} \\ \mathcal{L} & \xrightarrow{\tau} & \mathcal{L}^{\mathbb{C}}, \end{array}$$

where $\tilde{\tau}$ denotes the complexification $U \mapsto U \otimes \text{Id}$.

Correspondingly to [2], Proposition 1.3 we have

PROPOSITION A.1. *The mapping*

$$\rho^{\mathbb{C}} \circ \Phi : \mathcal{U}(\mathcal{H}) \longrightarrow \mathcal{L}^{\mathbb{C}}$$

is a homeomorphism, where Φ is defined in the same way as in [2], p. 7.

Now we must add the following lemma to [2], following Definition 1.4:

LEMMA A.2. *Let $\lambda \in \mathcal{L}$ and $\lambda = U(\lambda_0^\perp)$. Then we have that*

$$\lambda \in \mathcal{FL}_{\lambda_0}(\mathcal{H}) \iff U\bar{U}^{-1} + \text{Id} \text{ is a Fredholm operator.}$$

PROOF. By identifying $\mathcal{H} \cong \lambda_0 \otimes \mathbb{C} \cong \lambda_0 \oplus \sqrt{-1}\lambda_0$, we represent U as

$$U = X + \sqrt{-1}Y$$

with $X, Y : \lambda_0 \rightarrow \lambda_0$. Since

$$U\bar{U}^{-1} + \bar{U}\bar{U}^{-1} = (U + \bar{U})\bar{U}^{-1} = 2X\bar{U}^{-1},$$

we have that $U\bar{U}^{-1} + \text{Id}$ is a Fredholm operator, if and only if X is a Fredholm operator.

First we assume $\lambda \in \mathcal{FL}_{\lambda_0}(\mathcal{H})$. Then we know already by [2], Equation (1.1) that

$$(A.4) \quad \ker X = \lambda \cap \lambda_0,$$

and we have also

$$(A.5) \quad \lambda + \lambda_0 = \{-Y(x) + \sqrt{-1}X(x) + y \mid x, y \in \lambda_0\}.$$

So

$$(A.6) \quad \mathcal{H}/(\lambda + \lambda_0) \cong \lambda_0/\text{range}(X),$$

hence $\text{range}(X)$ is closed and of finite codimension in λ_0 .

Conversely, if X is a Fredholm operator, then also by (A.4), (A.5), and (A.6) we have $\lambda \in \mathcal{FL}_{\lambda_0}(\mathcal{H})$. \square

References

- [1] B. BOJARSKI, The abstract linear conjugation problem and Fredholm pairs of subspaces, in: "In Memoriam I.N. Vekua", Tbilisi Univ., Tbilisi, 1979, pp. 45–60 (Russian).
- [2] B. BOOSS-BAVNBEEK AND K. FURUTANI, The Maslov index – a functional analytical definition and the spectral flow formula, *Tokyo J. Math.* **21** (1998), 1–34.
- [3] —, —, Symplectic functional analysis and spectral invariants, in: B. Booss-Bavnbek, K.P. Wojciechowski (eds.), "Geometric Aspects of Partial Differential Equations", Amer. Math. Soc. Series *Contemporary Mathematics*, vol. 242, Providence, R.I., 1999, pp. 53–83.
- [4] B. BOOSS-BAVNBEEK AND K. P. WOJCIECHOWSKI, "Elliptic Boundary Problems for Dirac Operators", Birkhäuser, Boston, 1993.
- [5] J. BRÜNING AND M. LESCH, Spectral theory of boundary value problems for Dirac type operators, in: B. Booss-Bavnbek, K.P. Wojciechowski (eds.), "Geometric Aspects of Partial Differential Equations", Amer. Math. Soc. Series *Contemporary Mathematics*, vol. 242, Providence, R.I., 1999, pp. 203–215.
- [6] S.E. CAPPELL, R. LEE, AND E.Y. MILLER, On the Maslov index, *Comm. Pure Appl. Math.* **47** (1994), 121–186.
- [7] —, —, —, Selfadjoint elliptic operators and manifold decompositions Part I: Low eigenmodes and stretching, *Comm. Pure Appl. Math.* **49** (1996), 825–866. Part II: Spectral flow and Maslov index, *Comm. Pure Appl. Math.* **49** (1996), 869–909. Part III: Determinant line bundles and Lagrangian intersection, *Comm. Pure Appl. Math.* **52** (1999), 543–611.
- [8] M. DANIEL AND P. KIRK, WITH AN APPENDIX BY K.P. WOJCIECHOWSKI, "A general splitting formula for the spectral flow", Bloomington preprint, 1999.
- [9] M. DE GOSSON, The structure of q -symplectic geometry, *J. Math. Pures Appl.* **71** (1992), 429–453.
- [10] —, "Maslov Classes, Metaplectic Representation and Lagrangian Quantization", Mathematical Research vol. **95**, Akademie Verlag, Berlin 1997.
- [11] M. LESCH AND N. PEYERIMHOFF, On index formulas for manifolds with metric horns, *Comm. Partial Diff. Equations* **23** (1998), 649–684.
- [12] L. NICOLAESCU, The Maslov index, the spectral flow, and decomposition of manifolds, *Duke Math. J.* **80** (1995), 485–533.
- [13] —, "Generalized Symplectic Geometries and the Index of Families of Elliptic Problems", Mem. Amer. Math. Soc. vol. 609, Providence, 1997.
- [14] N. OTSUKI AND K. FURUTANI, Spectral flow and Maslov index arising from Lagrangian intersections, *Tokyo J. Math.* **14** (1991), 135–150.
- [15] J. PHILLIPS, Self-adjoint Fredholm operators and spectral flow, *Canad. Math. Bull.* **39** (1996), 460–467.
- [16] T. YOSHIDA, Floer homology and splittings of manifolds, *Ann. of Math.* **134** (1991), 277–323.

INSTITUT FOR MATEMATIK OG FYSIK, ROSKILDE UNIVERSITY, DK-4000 ROSKILDE, DENMARK

E-mail address: booss@mmf.ruc.dk

DEPARTMENT OF MATHEMATICS, SCIENCE UNIVERSITY OF TOKYO, NODA-SHI, CHIBA-KEN 278 JAPAN

E-mail address: furutani@ma.noda.sut.ac.jp

DEPARTMENT OF MATHEMATICS, SCIENCE UNIVERSITY OF TOKYO, NODA-SHI, CHIBA-KEN 278 JAPAN

E-mail address: nobukazu@ma.noda.sut.ac.jp

Liste over tidligere udkomne tekster
tilsendes gerne. Henvendelse herom kan
ske til IMFUFA's sekretariat

tlf. 46 74 22 63

-
- 217/92 "Two papers on APPLICATIONS AND MODELLING
IN THE MATHEMATICS CURRICULUM"
by: Mogens Niss
- 218/92 "A Three-Square Theorem"
by: Lars Kadison
- 219/92 "RUPNOK - stationær strømning i elastiske rør"
af: Anja Boisen, Karen Birkelund, Mette Olufsen
Vejleder: Jesper Larsen
- 220/92 "Automatisk diagnosticering i digitale kredsløb"
af: Bjørn Christensen, Ole Møller Nielsen
Vejleder: Stig Andur Pedersen
- 221/92 "A BUNDLE VALUED RADON TRANSFORM, WITH
APPLICATIONS TO INVARIANT WAVE EQUATIONS"
by: Thomas P. Branson, Gestur Olafsson and
Henrik Schlichtkrull
- 222/92 On the Representations of some Infinite Dimensional
Groups and Algebras Related to Quantum Physics
by: Johnny T. Ottesen
- 223/92 THE FUNCTIONAL DETERMINANT
by: Thomas P. Branson
- 224/92 UNIVERSAL AC CONDUCTIVITY OF NON-METALLIC SOLIDS AT
LOW TEMPERATURES
by: Jeppe C. Dyre
- 225/92 "HATMODELLEN" Impedansspektroskopi i ultrarent
en-krystallinsk silicium
af: Anja Boisen, Anders Gorm Larsen, Jesper Varmer,
Johannes K. Nielsen, Kit R. Hansen, Peter Bøggild
og Thomas Hougaard
Vejleder: Petr Viscor
- 226/92 "METHODS AND MODELS FOR ESTIMATING THE GLOBAL
CIRCULATION OF SELECTED EMISSIONS FROM ENERGY
CONVERSION"
by: Bent Sørensen

- 227/92 "Computersimulering og fysik"
af: Per M. Hansen, Steffen Holm,
Peter Maibom, Mads K. Dall Petersen,
Pernille Postgaard, Thomas B. Schrøder,
Ivar P. Zeck
Vejleder: Peder Voetmann Christiansen

- 228/92 "Teknologi og historie"
Fire artikler af:
Mogens Niss, Jens Høyrup, Ib Thiersen,
Hans Heddal

- 229/92 "Masser af information uden betydning"
En diskussion af informationsteorien
i Tor Nørretranders' "Mærk Verden" og
en skitse til et alternativ baseret
på andenordens kybernetik og semiotik.
af: Søren Brier

- 230/92 "Vinklens tredeling - et klassisk
problem"
et matematisk projekt af
Karen Birkelund, Bjørn Christensen
Vejleder: Johnny Ottesen

- 231A/92 "Elektrondiffusion i silicium - en
matematisk model"
af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen
Vejledere: Johnny Ottesen, H.B. Hansen

- 231B/92 "Elektrondiffusion i silicium - en
matematisk model" Kildetekster
af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen
Vejledere: Johnny Ottesen, H.B. Hansen

- 232/92 "Undersøgelse om den simultane opdagelse
af energiens bevarelse og isærdeles om
de af Mayer, Colding, Joule og Helmholtz
udførte arbejder"
af: L. Arleth, G.I. Dybkjær, M.T. Østergård
Vejleder: Dorte Posselt

- 233/92 "The effect of age-dependent host
mortality on the dynamics of an endemic
disease and
Instability in an SIR-model with age-
dependent susceptibility
by: Viggo Andreassen

- 234/92 "THE FUNCTIONAL DETERMINANT OF A FOUR-DIMENSIONAL
BOUNDARY VALUE PROBLEM"
by: Thomas P. Branson and Peter B. Gilkey

- 235/92 OVERFLADESTRUKTUR OG POREUDVIKLING AF KOKS
- Modul 3 fysik projekt -
af: Thomas Jessen
-

- 236a/93 INTRODUKTION TIL KVANTE
HALL EFFEKTEN
af: Anja Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 236b/93 STRØMSSAMMENBRUD AF KVANTE
HALL EFFEKTEN
af: Anja Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 237/93 The Wedderburn principal theorem and
Shukla cohomology
af: Lars Kadison
- 238/93 SEMIOTIK OG SYSTEMEGENSKABER (2)
Vektorbånd og tensorer
af: Peder Voetmann Christiansen
- 239/93 Valgsystemer - Modelbygning og analyse
Matematik 2. modul
af: Charlotte Gjerrild, Jane Hansen,
Maria Hermannsson, Allan Jørgensen,
Ragna Clauson-Kaas, Poul Lützen
Vejleder: Mogens Niss
- 240/93 Patologiske eksempler.
Om sære matematiske fisks betydning for
den matematiske udvikling
af: Claus Dråby, Jørn Skov Hansen, Runa
Ulsøe Johansen, Peter Meibom, Johannes
Kristoffer Nielsen
Vejleder: Mogens Niss
- 241/93 FOTOVOLTAISK STATUSNOTAT 1
af: Bent Sørensen
- 242/93 Brovedligeholdelse - bevar mig vel
Analyse af vejdirektoratets model for
optimering af broreparationer
af: Linda Kyndlev, Kare Fundal, Kamma
Tulinus, Ivar Zeck
Vejleder: Jesper Larsen
- 243/93 TANKEEKSPERIMENTER I FYSIKKEN
Et 1.modul fysikprojekt
af: Karen Birkelund, Stine Sofia Korremann
Vejleder: Dorthe Posselt
- 244/93 RADONTRANSFORMATIONEN og dens anvendelse
i CT-scanning
Projektrapport
af: Trine Andreasen, Tine Guldager Christiansen,
Nina Skov Hansen og Christine Iversen
Vejledere: Gestur Olafsson og Jesper Larsen
- 245a+b
/93 Time-Of-Flight målinger på krystallinske
halvledere
Specialerapport
af: Linda Szkotak Jensen og Lise Odgaard Gade
Vejledere: Petr Viscor og Niels Boye Olsen
- 246/93 HVERDAGSVIDEN OG MATEMATIK
- LÆREPROCESSER I SKOLEN
af: Lena Lindenskov, Statens Humanistiske
Forskningsråd, RUC, IMFUFA
- 247/93 UNIVERSAL LOW TEMPERATURE AC CON-
DUCTIVITY OF MACROSCOPICALLY
DISORDERED NON-METALS
by: Jeppe C. Dyre
- 248/93 DIRAC OPERATORS AND MANIFOLDS WITH
BOUNDARY
by: B. Booss-Bavnbek, K.P.Wojciechowski
- 249/93 Perspectives on Teichmüller and the
Jahresbericht Addendum to Schappacher,
Scholz, et al.
by: B. Booss-Bavnbek
With comments by W.Abikoff, L.Ahlfors,
J.Cerf, P.J.Davis, W.Fuchs, F.P.Gardiner,
J.Jost, J.-P.Kahane, R.Lohan, L.Lorch,
J.Radkau and T.Söderqvist
- 250/93 EULER OG BOLZANO - MATEMATISK ANALYSE SET I ET
VIDENSKABSTEORETISK PERSPEKTIV
Projektrapport af: Anja Juul, Lone Michelsen,
Tomas Højgård Jensen
Vejleder: Stig Andur Pedersen
- 251/93 Genotypic Proportions in Hybrid Zones
by: Freddy Bugge Christiansen, Viggo Andreasen
and Ebbe Thue Poulsen
- 252/93 MODELLERING AF TILFELDIGE FÆNOMENER
Projektrapport af: Birthe Friis, Lisbeth Helmgård
Kristina Charlotte Jakobsen, Marina Mosbek
Johannessen, Lotte Ludvigsen, Mette Hase Nielsen
- 253/93 Kuglepakning
Teori og model
af: Lise Arleth, Kåre Fundal, Nils Kruse
Vejleder: Mogens Niss
- 254/93 Regressionsanalyse
Materiale til et statistikkursus
af: Jørgen Larsen
- 255/93 TID & BETINGET UAFHÆNGIGHED
af: Peter Harremoës
- 256/93 Determination of the Frequency Dependent
Bulk Modulus of Liquids Using a Piezo-
electric Spherical Shell (Preprint)
by: T. Christensen and N.B.Olsen
- 257/93 Modellering af dispersion i piezoelektriske
keramikker
af: Pernille Postgaard, Jannik Rasmussen,
Christina Specht, Mikko Østergård
Vejleder: Tage Christensen
- 258/93 Supplerende kursusrmateriale til
"Lineære strukturer fra algebra og analyse"
af: Mogens Brun Heefelt
- 259/93 STUDIES OF AC HOPPING CONDUCTION AT LOW
TEMPERATURES
by: Jeppe C. Dyre
- 260/93 PARTITIONED MANIFOLDS AND INVARIANTS IN
DIMENSIONS 2, 3, AND 4
by: B. Booss-Bavnbek, K.P.Wojciechowski

- 261/93 OPGAVESAMLING
Bredde-kursus i Fysik
Eksamensopgaver fra 1976-93
- 262/93 Separability and the Jones Polynomial
by: Lars Kadison
- 263/93 Supplerende kursusmateriale til
"Lineære strukturer fra algebra
og analyse" II
af: Mogens Brun Heefelt
- 264/93 FOTOVOLTAISK STATUSNOTAT 2
af: Bent Sørensen
-
- 265/94 SPHERICAL FUNCTIONS ON ORDERED
SYMMETRIC SPACES
To Sigurdur Helgason on his
sixtyfifth birthday
by: Jacques Faraut, Joachim Hilgert
and Gestur Olafsson
- 266/94 Kommensurabilitets-oscillationer i
laterale supergitre
Fysikspeciale af: Anja Boisen,
Peter Bøggild, Karen Birkelund
Vejledere: Rafael Taboryski, Poul Erik
Lindelof, Peder Voetmann Christiansen
- 267/94 Kom til kort med matematik på
Eksperimentarium - Et forslag til en
opstilling
af: Charlotte Gjerrild, Jane Hansen
Vejleder: Bernhelm Booss-Bavnbek
- 268/94 Life is like a sewer ...
Et projekt om modellering af aorta via
en model for strømning i kloakrør
af: Anders Marcussen, Anne C. Nilsson,
Lone Michelsen, Per M. Hansen
Vejleder: Jesper Larsen
- 269/94 Dimensionsanalyse en introduktion
metaprojekt, fysik
af: Tine Guldager Christiansen,
Ken Andersen, Nikolaj Hermann,
Jannik Rasmussen
Vejleder: Jens Højgaard Jensen
- 270/94 THE IMAGE OF THE ENVELOPING ALGEBRA
AND IRREDUCIBILITY OF INDUCED REPRESENTATIONS OF EXPONENTIAL LIE GROUPS
by: Jacob Jacobsen
- 271/94 Matematikken i Fysikken.
Opdaget eller opfundet
NAT-BAS-projekt
vejleder: Jens Højgaard Jensen
- 272/94 Tradition og fornyelse
Det praktiske elevarbejde i gymnasiets
fysikundervisning, 1907-1988
af: Kristian Hoppe og Jeppe Guldager
Vejledning: Karin Beyer og Nils Hybel
- 273/94 Model for kort- og mellemdistanceløb
Verifikation af model
af: Lise Fabricius Christensen, Helle Pilemann,
Bettina Sørensen
Vejleder: Mette Olufsen
- 274/94 MODEL 10 - en matematisk model af intravenøse
anæstetikas farmakokinetik
3. modul matematik, forår 1994
af: Trine Andreassen, Bjørn Christensen, Christine
Green, Anja Skjoldborg Hansen. Lisbeth
Helmgård
Vejledere: Viggo Andreassen & Jesper Larsen
- 275/94 Perspectives on Teichmüller and the Jahresbericht
2nd Edition
by: Bernhelm Booss-Bavnbek
- 276/94 Dispersionsmodellering
Projektrapport 1. modul
af: Gitte Andersen, Rehannah Borup, Lisbeth Friis,
Per Gregersen, Kristina Vejro
Vejleder: Bernhelm Booss-Bavnbek
- 277/94 PROJEKTARBEJDSPEDAGOGIK - Om tre tolkninger af
problemorienteret projektarbejde
af: Claus Flensted Behrens, Frederik Voetmann
Christiansen, Jørn Skov Hansen, Thomas
Thingstrup
Vejleder: Jens Højgaard Jensen
- 278/94 The Models Underlying the Anaesthesia
Simulator Sophus
by: Mette Olufsen(Math-Tech), Finn Nielsen
(RISØ National Laboratory), Per Føge Jensen
(Herlev University Hospital), Stig Andur
Pedersen (Roskilde University)
- 279/94 Description of a method of measuring the shear
modulus of supercooled liquids and a comparison
of their thermal and mechanical response
functions.
af: Tage Christensen
- 280/94 A Course in Projective Geometry
by Lars Kadison and Matthias T. Kromann
- 281/94 Modellering af Det Cardiovasculære System med
Neural Puls kontrol
Projektrapport udarbejdet af:
Stefan Frello, Runa Ulsøe Johansen,
Michael Poul Curt Hansen, Klaus Dahl Jensen
Vejleder: Viggo Andreassen
- 282/94 Parallelle algoritmer
af: Erwin Dan Nielsen, Jan Danielsen,
Niels Bo Johansen

- 283/94 Grænser for tilfældighed
(en kaotisk talgenerator)
af: Erwin Dan Nielsen og Niels Bo Johansen
- 284/94 Det er ikke til at se det, hvis man ikke
lige ve' det!
Gymnasie matematikkens begrundelsesproblem
En specialerapport af Peter Hauge Jensen
og Linda Kyndlev
Vejleder: Mogens Niss
- 285/94 Slow coevolution of a viral pathogen and
its diploid host
by: Viggo Andreassen and
Freddy B. Christiansen
- 286/94 The energy master equation: A low-temperature
approximation to Bässler's random walk model
by: Jeppe C. Dyre
- 287/94 A Statistical Mechanical Approximation for the
Calculation of Time Auto-Correlation Functions
by: Jeppe C. Dyre
- 288/95 PROGRESS IN WIND ENERGY UTILIZATION
by: Bent Sørensen
- 289/95 Universal Time-Dependence of the Mean-Square
Displacement in Extremely Rugged Energy
Landscapes with Equal Minima
by: Jeppe C. Dyre and Jacob Jacobsen
- 290/95 Modellering af uregelmæssige bølger
Et 3.modul matematik projekt
af: Anders Marcussen, Anne Charlotte Nilsson,
Lone Michelsen, Per Mørkegaard Hansen
Vejleder: Jesper Larsen
- 291/95 1st Annual Report from the project
LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH
ENERGY SYSTEM
an example of using methods developed for the
OECD/IEA and the US/EU fuel cycle externality study
by: Bent Sørensen
- 292/95 Fotovoltaisk Statusnotat 3
af: Bent Sørensen
- 293/95 Geometridiskussionen - hvor blev den af?
af: Lotte Ludvigsen & Jens Frandsen
Vejleder: Anders Madsen
- 294/95 Universets udvidelse -
et metaprojekt
Af: Jesper Duelund og Birthe Friis
Vejleder: Ib Lundgaard Rasmussen
- 295/95 A Review of Mathematical Modeling of the
Controlled Cardiovascular System
By: Johnny T. Ottesen
- 296/95 RETIKULER den klassiske mekanik
af: Peder Voetmann Christiansen
- 297/95 A fluid-dynamical model of the aorta with
bifurcations
by: Mette Olufsen and Johnny Ottesen
- 298/95 Mordet på Schrödingers kat - et metaprojekt om
to fortolkninger af kvantemekanikken
af: Maria Hermannsson, Sebastian Horst,
Christina Specht
Vejledere: Jeppe Dyre og Peder Voetmann Christiansen
- 299/95 ADAM under figenbladet - et kig på en samfunds-
videnskabelig matematisk model
Et matematisk modelprojekt
af: Claus Dræby, Michael Hansen, Tomas Højgård Jensen
Vejleder: Jørgen Larsen
- 300/95 Scenarios for Greenhouse Warming Mitigation
by: Bent Sørensen
- 301/95 TOK Modellering af træers vækst under påvirkning
af ozon
af: Glenn Møller-Holst, Marina Johannessen, Birthe
Nielsen og Bettina Sørensen
Vejleder: Jesper Larsen
- 302/95 KOMPRESSORER - Analyse af en matematisk model for
aksialkompressor
Projektrapport af: Stine Bøggild, Jakob Hilmer,
Pernille Postgaard
Vejleder: Viggo Andreassen
- 303/95 Masterlignings-modeller af Glasovergangen
Termisk-Mekanisk Relaksation
Specialerapport udarbejdet af:
Johannes K. Nielsen, Klaus Dahl Jensen
Vejledere: Jeppe C. Dyre, Jørgen Larsen
- 304a/95 STATISTIKNOTER Simple binomialfordelingsmodeller
af: Jørgen Larsen
- 304b/95 STATISTIKNOTER Simple normalfordelingsmodeller
af: Jørgen Larsen
- 304c/95 STATISTIKNOTER Simple Poissonfordelingsmodeller
af: Jørgen Larsen
- 304d/95 STATISTIKNOTER Simple multinomialfordelingsmodeller
af: Jørgen Larsen
- 304e/95 STATISTIKNOTER Mindre matematisk-statistisk opslagsværk
indeholdende bl.a. ordforklaringer, resuméer og
tabeller
af: Jørgen Larsen

- 305/95 The Maslov Index:
A Functional Analytical Definition
And The Spectral Flow Formula
By: B. Booss-Bavnbek, K. Furutani
- 306/95 Goals of mathematics teaching
Preprint of a chapter for the forthcoming International Handbook of Mathematics Education (Alan J. Bishop, ed)
By: Mogens Niss
- 307/95 Habit Formation and the Thirdness of Signs
Presented at the semiotic symposium
The Emergence of Codes and Intensions as a Basis of Sign Processes
By: Peder Voetmann Christiansen
- 308/95 Metaforer i Fysikken
af: Marianne Wilcken Bjerregaard, Frederik Voetmann Christiansen, Jørn Skov Hansen, Klaus Dahl Jensen, Ole Schmidt
Vejledere: Peder Voetmann Christiansen og Petr Viscor
- 309/95 Tiden og Tanken
En undersøgelse af begrebsverdenen Matematik udført ved hjælp af en analogi med tid
af: Anita Stark og Randi Petersen
Vejleder: Bernhelm Booss-Bavnbek
-
- 310/96 Kursusmateriale til "Lineære strukturer fra algebra og analyse" (E1)
af: Mogens Brun Heefelt
- 311/96 2nd Annual Report from the project
LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH ENERGY SYSTEM
by: Hélène Connor-Lajambe, Bernd Kuemmel, Stefan Krüger Nielsen, Bent Sørensen
- 312/96 Grassmannian and Chiral Anomaly
by: B. Booss-Bavnbek, K.P. Wojciechowski
- 313/96 THE IRREDUCIBILITY OF CHANCE AND THE OPENNESS OF THE FUTURE
The Logical Function of Idealism in Peirce's Philosophy of Nature
By: Helmut Pape, University of Hannover
- 314/96 Feedback Regulation of Mammalian Cardiovascular System
By: Johnny T. Ottesen
- 315/96 "Rejsen til tidens indre" - Udarbejdelse af a + b et manuskript til en fjernsynsudsendelse + manuskript
af: Gunhild Hune og Karina Goyle
Vejledere: Peder Voetmann Christiansen og Bruno Ingemann
- 316/96 Plasmaoscillation i natriumklynger
Specialerapport af: Peter Meibom, Mikko Østergård
Vejledere: Jeppe Dyre & Jørn Borggreen
- 317/96 Poincaré og symplektiske algoritmer
af: Ulla Rasmussen
Vejleder: Anders Madsen
- 318/96 Modelling the Respiratory System
by: Tine Guldager Christiansen, Claus Dræby
Supervisors: Viggo Andreassen, Michael Danielsen
- 319/96 Externality Estimation of Greenhouse Warming Impacts
by: Bent Sørensen
- 320/96 Grassmannian and Boundary Contribution to the -Determinant
by: K.P. Wojciechowski et al.
- 321/96 Modelkompetencer - udvikling og afprøvning af et begrebsapparat
Specialerapport af: Nina Skov Hansen, Christine Iversen, Kristin Troels-Smith
Vejleder: Morten Blomhøj
- 322/96 OPGAVESAMLING
Bredde-Kursus i Fysik 1976 - 1996
- 323/96 Structure and Dynamics of Symmetric Diblock Copolymers
PhD Thesis
by: Christine Maria Papadakis
- 324/96 Non-linearity of Baroreceptor Nerves
by: Johnny T. Ottesen
- 325/96 Retorik eller realitet ?
Anvendelser af matematik i det danske Gymnasiums matematikundervisning i perioden 1903 - 88
Specialerapport af Helle Pilemann
Vejleder: Mogens Niss
- 326/96 Bevisteorier
Eksemplificeret ved Gentzens bevis for konsistensen af teorien om de naturlige tal
af: Gitte Andersen, Lise Mariane Jeppesen, Klaus Frovin Jørgensen, Ivar Peter Zeck
Vejledere: Bernhelm Booss-Bavnbek og Stig Andur Pedersen
- 327/96 NON-LINEAR MODELLING OF INTEGRATED ENERGY SUPPLY AND DEMAND MATCHING SYSTEMS
by: Bent Sørensen
- 328/96 Calculating Fuel Transport Emissions
by: Bernd Kuemmel

- 329/96 The dynamics of cocirculating influenza strains conferring partial cross-immunity and
A model of influenza A drift evolution
by: Viggo Andreasen, Juan Lin and Simon Levin
- 330/96 LONG-TERM INTEGRATION OF PHOTOVOLTAICS INTO THE GLOBAL ENERGY SYSTEM
by: Bent Sørensen
- 331/96 Viskøse fingre
Specialerapport af:
Vibeke Orlén og Christina Specht
Vejledere: Jacob M. Jacobsen og Jesper Larsen
-
- 332/97 ANOMAL SWELLING AF LIPIDE DOBBELTLAG
Specialerapport af:
Stine Sofia Korremann
Vejleder: Dorthe Posselt
- 333/97 Biodiversity Matters
an extension of methods found in the literature on monetisation of biodiversity
by: Bernd Kuemmel
- 334/97 LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH ENERGY SYSTEM
by: Bernd Kuemmel and Bent Sørensen
- 335/97 Dynamics of Amorphous Solids and Viscous Liquids
by: Jeppe C. Dyre
- 336/97 PROBLEM-ORIENTATED GROUP PROJECT WORK AT ROSKILDE UNIVERSITY
by: Kathrine Legge
- 337/97 Verdensbankens globale befolkningsprognose - et projekt om matematisk modellering
af: Jørn Chr. Bendtsen, Kurt Jensen, Per Pauli Petersen
Vejleder: Jørgen Larsen
- 338/97 Kvantisering af nanolederes elektriske ledningsevne
Første modul fysikprojekt
af: Søren Dam, Esben Danielsen, Martin Niss, Esben Friis Pedersen, Frederik Resen Steenstrup
Vejleder: Tage Christensen
- 339/97 Defining Discipline
by: Wolfgang Coy
- 340/97 Prime ends revisited - a geometric point of view -
by: Carsten Lunde Petersen
- 341/97 Two chapters on the teaching, learning and assessment of geometry
by Mogens Niss
- 342/97 LONG-TERM SCENARIOS FOR GLOBAL ENERGY DEMAND AND SUPPLY
A global clean fossil scenario discussion paper prepared by Bernd Kuemmel
Project leader: Bent Sørensen
- 343/97 IMPORT/EKSPORT-POLITIK SOM REDSKAB TIL OPTIMERET UDNYTTELSE AF EL PRODUCERET PÅ VE-ANLÆG
af: Peter Meibom, Torben Svendsen, Bent Sørensen
- 344/97 Puzzles and Siegel disks
by Carsten Lunde Petersen
-
- 345/98 Modeling the Arterial System with Reference to an Anesthesia Simulator
Ph.D. Thesis
by: Mette Sofie Olufsen
- 346/98 Klyngedannelse i en hulkatode-forstøvningsproces
af: Sebastian Horst
Vejledere: Jørn Borggren, NBI, Niels Boye Olsen
- 347/98 Verificering af Matematiske Modeller - en analyse af Den Danske Eulerske Model
af: Jonas Blomqvist, Tom Pedersen, Karen Timmermann, Lisbet Øhlenschläger
Vejleder: Bernhelm Booss-Bavnbek
- 348/98 Case study of the environmental permission procedure and the environmental impact assessment for power plants in Denmark
by: Stefan Krüger Nielsen
Project leader: Bent Sørensen
- 349/98 Tre rapporter fra FAGMAT - et projekt om tal og faglig matematik i arbejdsmarkedsuddannelserne
af: Lena Lindenskov og Tine Wedege
- 350/98 OPGAVERSAMLING - Bredde-Kursus i Fysik 1976 - 1998
Erstatter teksterne 3/78, 261/93 og 322/96
- 351/98 Aspects of the Nature and State of Research in Mathematics Education
by: Mogens Niss

- 352/98 The Herman-Swiatec Theorem with applications
by: Carsten Lunde Petersen
- 353/98 Problemløsning og modellering i en almindelig matematikundervisning
Specialerapport af: Per Gregersen og Tomas Højgaard Jensen
Vejleder: Morten Blomhøj
- 354/98 A GLOBAL RENEWABLE ENERGY SCENARIO
by: Bent Sørensen and Peter Meibom
- 355/98 Convergence of rational rays in parameter spaces
by: Carsten Lunde Petersen and Gustav Ryd
- 356/98 Terrænmodellering
Analyse af en matematisk model til konstruktion af terrænmodeller
Modelprojekt af: Thomas Frommelt, Hans Ravnkjær Larsen og Arnold Skimminge
Vejleder: Johnny Ottesen
- 357/98 Cayleys Problem
En historisk analyse af arbejdet med Cayley problem fra 1870 til 1918
Et matematisk videnskabsfagsprojekt af: Rikke Degn, Bo Jakobsen, Bjarke K.W. Hansen, Jesper S. Hansen, Jesper Udesen, Peter C. Wulff
Vejleder: Jesper Larsen
- 358/98 Modeling of Feedback Mechanisms which Control the Heart Function in a View to an Implementation in Cardiovascular Models
Ph.D. Thesis by: Michael Danielsen
- 359/99 Long-Term Scenarios for Global Energy Demand and Supply Four Global Greenhouse Mitigation Scenarios
by: Bent Sørensen
- 360/99 SYMMETRI I FYSIK
En Meta-projektrapport af: Martin Niss, Bo Jakobsen & Tine Bjarke Bonné
Vejleder: Peder Voetmann Christiansen
- 361/99 Symplectic Functional Analysis and Spectral Invariants
by: Bernhelm Booss-Bavnbek, Kenro Furutani
- 362/99 Er matematik en naturvidenskab? - en udspejling af diskussionen
En videnskabsfagsprojekt-rapport af Martin Niss
Vejleder: Mogens Nørgaard Olesen
- 363/99 EMERGENCE AND DOWNWARD CAUSATION
by: Donald T. Campbell, Mark H. Bickhard and Peder V. Christiansen
- 364/99 Illustrationens kraft
Visuel formidling af fysik
Integreret speciale i fysik og kommunikation
af: Sebastian Horst
Vejledere: Karin Beyer, Søren Kjørup
- 365/99 To know - or not to know - mathematics, that is a question of context
by: Tine Wedege
- 366/99 LATEX FOR FORFATTERE
En introduktion til LATEX og IMFUFA-LATEX
af: Jørgen Larsen
- 367/99 Boundary Reduction of Spectral invariants and Unique Continuation Property
by Bernhelm Booss-Bavnbek
- 368/99 Kvartalsrapport for projektet
SCENARIER FOR SAMLET UDNYTTELSE AF BRINT SOM ENERGIBÆRER I DANMARKS FREMTIDIGE ENERGISTYSTEM
Projektleder: Bent Sørensen
- 369/99 DYNAMICS OF Complex Quadratic Correspondences
by: Jacob Jalving
- 370/99 OPGAVESAMLING
Bredde-Kursus i Fysik 1976 - 1999
(erstatte tekst nr. 350/98)
- 370/99 Bevisets stilling
- beviser og bevisførelse i en gymnasial matematikundervisning
Matematikspeciale af: Maria Hermannsson
Vejleder: Mogens Niss